

The Cardinality of the Second Uniform Indiscernible

Greg Hjorth

Group in Logic, University of California, Berkeley, CA94720

February 1, 2008

Abstract

When the second uniform indiscernible is $\aleph_2$, the Martin-Solovay tree only constructs countably many reals; this resolves a number of open questions in descriptive set theory.¹

1. Introduction and Definitions.

From now on we work in the theory $\text{ZFC} + \forall x \in \omega^\omega (x^\# \exists)$.

Assuming $\forall x \in \omega^\omega (x^\# \exists)$, Martin and Solovay showed that every Σ_3^1 set is the projection of a simply definable tree, T_2 . This extended several earlier results, but in some instances only with the further assumption that T_2 has size less than $\aleph_2$.

The present paper smooths the way for a further analysis by showing that if $u_2 = \aleph_2$ (a necessary and sufficient condition for T_2 to have size bigger than $\aleph_1$) then the smallest inner model of set theory containing T_2 and all the ordinals, $\mathbf{L}[T_2]$, has only countably many reals.

1.1. Definition: $\gamma \in \mathbf{Ord}$ is a *uniform indiscernible* if $\forall x \in \omega^\omega (\gamma \text{ is an } \mathbf{L}[x] \text{ indiscernible})$.

1.2. Notation: For $\alpha \in \mathbf{Ord}$, let u_α be the α^{th} uniform indiscernible, beginning with $u_1 = \aleph_1$. For $x \in \omega^\omega$, $\tau \in \mathcal{L}(\mathbf{L}[x])$ indicates that τ is a skolem function over $\mathbf{L}[x]$, definable from no parameters.

¹1991 *Mathematics Subject Classification*. Primary 04A15. *Key words and phrases*. Descriptive set theory, uniform indiscernibles, Martin-Solovay tree.

1.3.Representation Lemma (Solovay): For all $\gamma \in \text{Ord}$, $\beta < u_\gamma$, there exists $x \in \omega^\omega$, $\tau \in \mathcal{L}(\mathbf{L}[x])$, and $\gamma_1, \dots, \gamma_n < \gamma$ such that $\tau(u_{\gamma_1}, \dots, u_{\gamma_n}) = \beta$. $\square$

A similar result holds for finite strings of ordinals; any finite string of ordinals less than u_ω can be coded from a single real and finitely many of the uniform indiscernibles less than u_ω . These lemmas are significant because they enable us to phrase questions about what occurs inside $\mathbf{L}[x]$ at $\alpha < u_\omega$ in a $\Delta_3^1(x, y)$ manner, for any $y \in \omega^\omega$ from which α can be defined using the uniform indiscernibles.

For our purposes it will be unimportant how the Martin-Solovay tree is defined. It can, for example, be extracted from the scale discussed in Moschovakis [3]. A closely related tree arises from the “shift maps”, as implicit in the construction in 2..1. below. Both derive from [2]. The only properties of the tree required are that:

- (i) it is no worse than Σ_3^1 in the codes for the ordinals less than u_ω ;
- (ii) it projects to the complete Σ_3^1 set;
- (iii) it has the same cardinality as u_2 .

Let us fix such a tree and call it T_2 .

1.4.Notation: For $\alpha \in \mathbf{Ord}$, $x \in \omega^\omega$, set $\text{Next}(\alpha, x)$ to be the least $\mathbf{L}[x]$ indiscernible above α .

2. $u_2 = \aleph_2$.

2.1.Theorem($\text{ZFC} + \forall x \in \omega^\omega (x^\# \exists)$): If $u_2 = \aleph_2$, then there are only countably many reals in $\mathbf{L}[T_2]$.

Proof:

Suppose otherwise, and now we will derive a contradiction. Let θ be a big, regular cardinal, so that $\mathbf{V}_\theta \models \text{“}T_2 \text{ constructs } \geq \aleph_1 \text{ many reals”}$. Now choose

- (i) $N_0 \prec N_1 \prec \dots N_i \dots \prec \mathbf{V}_\theta$, such that $\forall i \in \omega$, $\aleph_1 \subset N_i$, $\|N_i\| = \aleph_1$;
- (ii) $(y_i)_{i \in \omega} \subset \omega^\omega$, $y_i \in N_{i+1}$, $y_i^\#$ recursive in y_{i+1} , such that $\text{Next}(\aleph_1, y_i) > \aleph_2 \cap N_i$.

Let $y \in \omega^\omega$ uniformly code the sequence $(y_i)_{i \in \omega}$, let $N_\omega = \bigcup (N_i)_{i \in \omega}$, let N be the transitive collapse of N_ω , and let T_2^* be T_2 as calculated in N . N must be correct in calculating that T_2^* constructs at least $\aleph_1$ many reals. So it will suffice to show that $T_2^* \in \mathbf{L}[y]$. Let $(w_\alpha)_{\alpha \in \omega_1}$ be a generic sequence of reals for $\text{Coll}(\omega, < \omega_1)$. In virtue of general facts about forcing, it will suffice to show that T_2^* is in $\mathbf{L}[y, (w_\alpha)_{\alpha \in \omega_1}]$. Since $(T_2)^{N[(w_\alpha)_{\alpha \in \omega_1}]} = (T_2)^N = T_2^*$ and since every ordinal less than $(u_\omega)^N$ is coded by some $(u_1)^N, \dots, (u_n)^N$

and $(w_{\alpha_1}, \dots, w_{\alpha_n}, y_0, \dots, y_n)$, it suffices to show that $\text{Th}_{\Sigma_3^1}^{N[(w_\alpha)_{\alpha \in \omega_1}]}((w_\alpha)_{\alpha \in \omega_1} \cup (y_i)_{i \in \omega})$ can be calculated by $\mathbf{L}[y, (w_\alpha)_{\alpha \in \omega_1}]$.

Let S be the set of Σ_3^1 sentences such that “ $\exists z \psi(w_{\alpha_1}, \dots, w_{\alpha_n}, y_0, \dots, y_n, z)$ ” $\in S$ provided:

- (i) ψ is Π_2^1 ;
- (ii) there exists x recursively above $(w_{\alpha_1}, \dots, w_{\alpha_n}, y_0, \dots, y_n)$, and a well-founded model $m(x) \models \exists z \psi(w_{\alpha_1}, \dots) \wedge \mathbf{V} = \mathbf{L}[x]$, along with indiscernibles $(c_i)_{i \in \omega, i > 0}$ which generate $m(x)$;
- (iii) and there exists $f : \mathbf{Ord}^{m(x)} \rightarrow (u_\omega)^N$, such that $m(x) \models \gamma = \tau(c_1, \dots) \in [c_m, c_{m+1})$ iff $f(\gamma) \in [(u_m)^N, (u_{m+1})^N)$, for all $\tau \in \mathcal{L}(M(x))$, and such that $f(\tau(c_1, \dots, c_n)) = \sigma((u_1)^N, \dots, (u_m)^N)$ implies that $f(\tau(c_{l_1}, \dots, c_{l_n})) = \sigma((u_{l_1})^N, \dots, (u_{l_m})^N)$, for all $\tau \in \mathcal{L}(m(x))$, $\sigma \in \mathcal{L}(\mathbf{L}[r])$, $r \in (\omega^\omega)^{N[(w_\alpha)_{\alpha \in \omega_1}]}$, $m < n$, $0 < l_1 < l_2, \dots, < l_n < \omega$.

Observe that membership in S can be phrased in terms of a tree construction, with a given f and $m(x)$ corresponding to a branch. We can view the nodes of the tree as constructing larger and larger finite initial segments of the theory of $m(x)[(c_i)_{0 < i < \omega}]$, as in a consistency property, along with larger and larger finite initial segments of f , that witnesses well foundedness in a particularly strong form. Observe that the tree can in fact be calculated in $\mathbf{L}[y, (w_\alpha)_{\alpha \in \omega_1}]$, since it can locate a set of reals $X \subset (\omega^\omega)^N$ which is closed under the pairing and sharp operations and provides codes for the ordinals below $(u_\omega)^N$. In essence, we may take X to be the set of reals generated by $(w_\alpha)_{\alpha \in \omega_1} \cup (y_i)_{i \in \omega}$ by the operations of pairing and taking sharps, and observe that the proof of Solovay’s representation lemma goes through for reals restricted to this set and for ordinals

Claim: if “ $\exists z \psi(w_{\alpha_1}, \dots)$ ” $\in S$, then $N[(w_\alpha)_{\alpha \in \omega_1}] \models \exists z \psi(w_{\alpha_1}, \dots)$.

Given a branch $(m(x), f)$, we can expand $m(x)$ out along ω_1 many indiscernibles, inducing a model $m(x)(c_\alpha)_{0 < \alpha < \omega_1}$. By Soenfield absoluteness, it suffices to show that this model is well founded. Now we witness well-foundedness by canonically extending f to $f_{\omega_1} : \mathbf{Ord}^{m(x)(c_\alpha)_{0 < \alpha < \omega_1}} \rightarrow (u_{\omega_1})^N$. Given $\tau(c_{\beta_1} \dots c_{\beta_n})$, where $\tau \in \mathcal{L}(m(x))$, we find $r \in (\omega^\omega)^N, \sigma \in \mathcal{L}(\mathbf{L}[r]), m \leq n$ such that $f(\tau(c_1 \dots c_n)) = \sigma((u_1)^N, \dots, (u_m)^N) \in [(u_m)^N, (u_{m+1})^N)$, and set $f_{\omega_1}(\tau(c_{\beta_1} \dots c_{\beta_n})) = \sigma((u_{\beta_1})^N \dots (u_{\beta_m})^N)$.

Subclaim: f_{ω_1} is well defined.

Suppose $\tau_1(c_{\beta_1} \dots c_{\beta_n}) = \tau_2(c_{\beta_k} \dots c_{\beta_n}), f(\tau_1(c_1 \dots c_n)) = \sigma_1((u_1)^N \dots (u_n)^N)$ and $f(\tau_2(c_1 \dots c_{n+1-k})) = \sigma_2((u_1)^N \dots (u_{n+1-k})^N)$ where σ_1 and σ_2 are skolem

functions definable from reals in N . Then since f respected the shift maps and the $(u_i)_{0 < i < \omega}^N$ are joint indiscernibles for the reals used in σ_1 and σ_2 , we have that $f(\tau_2(c_k \dots c_n)) = f(\tau_1(c_1 \dots c_n)) = \sigma_2((u_k)^N \dots (u_n)^N) = \sigma_1((u_1)^N \dots (u_n)^N)$, and, consequently, $\sigma_2((u_{\beta_k})^N \dots (u_{\beta_n})^N) = \sigma_1((u_{\beta_1})^N \dots (u_{\beta_n})^N)$, as required. The other cases follow in an exactly similar fashion.

Subclaim: f_{ω_1} is order preserving.

This is immediate given the previous claim and the requirement that f respect the order on $\mathbf{Ord}^{m(x)}$.

Claim: if $N[(w_\alpha)_{\alpha \in \omega_1}] \models \exists z \psi(w_{\alpha_1}, \dots)$ then “ $\exists z \psi(w_{\alpha_1}, \dots)$ ” ϵ S.

This follows by considering the appropriate sharp, and the natural map into $(u_\omega)^N$.

But since S can be calculated in $\mathbf{L}[y, (w_\alpha)_{\alpha \in \omega_1}]$, these last two claims suffice. $\square$

2.2. Corollaries (ZFC + $\forall x \in \omega^\omega (x^\sharp \exists)$):

- (i) every Σ_3^1 set of size greater than $\aleph_1$ contains a perfect set;
- (ii) Martin’s Axiom implies Σ_3^1 lebesgue measurability;
- (iii) if there is a Σ_3^1 well ordering of the reals, then the continuum hypothesis holds;
- (iv) if $\mathbf{K}_0 \subset \mathbf{V}$ is an inner model with a good Σ_3^1 well ordering of the reals, and if $u_2 = \aleph_2$, and $\mathbf{K}_0 \prec_{\Sigma_3^1} \mathbf{V}$, or the goodness of the well ordering is in some sense sufficiently robust, then $\mathbf{K}_0$ does not calculate ω_1 correctly.

Proof:

These results are all proved using standard techniques along with the fact that T_2 never constructs more than $\aleph_1$ many reals: if $u_2 = \aleph_2$, then this follows by the theorem; if $u_2 < \aleph_2$ then, after coding T_2 as subset of the ordinals, this follows by the entirely general observation that a subset of ω_1 never constructs more than $\aleph_1$ many reals. For instance, (iii) follows as in Kechris’s proof of Mansfield theorem that if there is a Σ_2^1 well ordering of the reals then every real is in $\mathbf{L}$. (ii) follows by considering that “almost every” real must be random over $\mathbf{L}[T_2]$. This answers a question of Judah’s. $\square$

2.3. Remark: 2.2.(iii) is rather strange sounding, but the point is that the standard canonical inner models of large cardinals below Π_2^1 determinacy have such “robust” well orders. This indicates difficulties in forcing models of $u_2 = \aleph_2$.

References

- [1] T. J. Jech, *Set theory*, Academic Press, London, New York, San Francisco, 1978.
- [2] D. A. Martin and R. M. Solovay, *A basis theorem for Σ_3^1 sets of reals*, Ann. of Math.(2) 89(1969), 138-160.
- [3] Y. N. Moscovakis, *Descriptive set theory*, North-Holland, Amsterdam, 1980.